

# Sabrex®corn

FERTILIZER GROUP 2 | Reg. No. B4659 Act No. 36 of 1947

Endophytic Trichoderma root inoculant for improved growth and productivity.

**Active ingredient:** 

Trichoderma spp..... min. 7 x 10<sup>7</sup> CFUs per gram

300g



Registered & Manufactured by:
Advanced Biological Marketing (Pty) Ltd | Co. Reg No: 2006/007975/07
PO Box 3253, Lichtenburg 2740, RSA
Tel No: 018 632 0362 | Email: info@abmafrica.co.za



# **DIRECTIONS FOR USE:** USE ONLY AS INDICATED

# SHAKE WELL BEFORE USE

- Apply as a dry seed treatment prior to planting.
- Sprinkle SabrEx® onto seed in bag.
- Treat in one bag increments to ensure uniform coating of seed with SabrEx®.
- Shake bag or mix until uniform seed coverage is obtained.
- Plant treated seed directly after SabrEx® seed treatment has been applied.

| TREATMENT      | DOSAGE / kg SEED  | COMMENT  |
|----------------|---|--|
| Seed Treatment | 300 g per 250 kg of seed<br>OR<br>1 Scoop (50 ml) per 1 bag (25 kg) of seed | Apply as a dry seed treatment at planting.   |
| In Furrow      | 30 g for every bag of seed per hectare                                      | Dilute with the appropriate amount of non-<br>chlorinated water and apply in furrow directly<br>on seed. |

# STORE IN A DRY PLACE BELOW 24°C & OUT OF DIRECT SUNLIGHT

## **IMPORTANT NOTES**

- Contents may settle. Shake well before use.
- Do not use inoculant after expiry date.

### **WARNINGS AND PRECAUTIONS:**

- Keep product out of direct sunlight.
- Do not inhale product dust.
- Avoid contact with eyes and skin. If skin or eye contact occurs, flush with copious amounts of clean water.
- Do not eat, drink or smoke while handling the product or before washing hands.
- Wash hands with soap and water after use.
- Destroy empty bag and do not use for any other purpose.
- Prevent contamination of food, feeds, drinking water and eating utensils.

Although this remedy has been extensively tested under a large variety of conditions, the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be affected by factors such as abnormal soil, climatic and storage conditions, quality of dilution water, compatibility with other substances not indicated on the label and the occurrence of resistance of the disease against the remedy concerned, as well as by the method, time and accuracy of application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment, or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in the event of any uncertainty.

